

Marco Antonio Borba da Silva

**Aspectos sobre a ecologia e taxonomia de fungos poliporoides  
(Basidiomycota) da Mata Atlântica de Santa Catarina, Brasil**

Dissertação submetida ao Programa de Pós-Graduação em Biologia de Fungos, Algas e Plantas da Universidade Federal de Santa Catarina para a obtenção do Grau de Mestre em Biologia de Fungos, Algas e Plantas.

Orientador: Prof. Dr. Elisandro Ricardo Drechsler dos Santos

Co-orientador: Prof. Dr. Gerardo Lucio Robledo

Florianópolis  
2013

Ficha de identificação da obra elaborada pelo autor,  
através do Programa de Geração Automática da Biblioteca Universitária da UFSC.

Borba-Silva, Marco Antonio  
Aspectos sobre a ecologia e taxonomia de fungos  
poliporoides (Basidiomycota) da Mata Atlântica de Santa  
Catarina, Brasil : Fungos poliporoides de Santa Catarina  
/ Marco Antonio Borba-Silva ; orientador, Elisandro  
Ricardo Drechsler dos Santos ; co-orientador, Gerardo  
Lucio Robledo. - Florianópolis, SC, 2013.  
68 p.

Dissertação (mestrado) - Universidade Federal de Santa  
Catarina, Centro de Ciências Biológicas. Programa de Pós-  
Graduação em Biologia de Fungos, Algas e Plantas.

Inclui referências

1. Biologia de Fungos, Algas e Plantas. 2. Ecologia de  
Fungos. 3. Taxonomia de Fungos. 4. Grupos funcionais. 5.  
Fungos poliporoides . I. Drechsler dos Santos, Elisandro  
Ricardo . II. Robledo, Gerardo Lucio. III. Universidade  
Federal de Santa Catarina. Programa de Pós-Graduação em  
Biologia de Fungos, Algas e Plantas. IV. Título.

Marco Antonio Borba da Silva

**Aspectos sobre a ecologia e taxonomia de fungos poliporoides  
(Basidiomycota) da Mata Atlântica de Santa Catarina, Brasil**

Esta Dissertação foi julgada adequada para obtenção do Título de “Mestre” e aprovada em sua forma final pelo Programa de Pós-Graduação em Biologia de Fungos, Algas e Plantas.

Local, 19 de abril de 2013.

---

Prof.<sup>a</sup> Dr.<sup>a</sup> Maria Alice Neves  
Coordenadora do Curso

**Banca Examinadora:**

---

Prof. Dr. Elisandro Ricardo Drechsler dos Santos,  
Orientador  
Universidade Federal de Santa Catarina

---

Prof. Dr. Carlos Urcelay,  
Universidad Nacional de Córdoba

---

Prof.<sup>a</sup> Dr.<sup>a</sup> Maria Alice Neves,  
Universidade Federal de Santa Catarina

---

Prof. Dr. Mateus A. Reck,  
Universidade Federal do Rio Grande do Sul



À minha Mãe, em memória;  
à Nayara, minha amada, dedico.



## AGRADECIMENTOS

Agradeço ao Prof. Dr. Elisandro Ricardo Drechsler dos Santos, meu orientador; por ter me dado a oportunidade de crescer profissionalmente, acreditando no meu potencial; pelas orientações, ensinamentos, palavras sinceras e amizade.

Agradeço ao Prof. Dr. Gerardo Lucio Robledo, meu co-orientador, pelo suporte técnico, orientação, paciência e amizade.

Agradeço a Prof<sup>ª</sup> Maria Alice Neves pela cooperação, ensinamentos e amizade.

Agradeço aos amigos e professores do Laboratório de Micologia – IMBIV – Universidad Nacional de Córdoba, não só pelo apoio técnico, mas principalmente por me acolherem como um amigo de longa data.

Agradeço aos meus amigos e colegas do Micolab pela convivência, cumplicidade, ajuda, amizade e pelas reuniões e saídas de campo.

Em especial, aos meus amigos micólogos Carlos Salvador Montoya, Diogo Costa Rezende, Fernando Mafalda Freire, Salomé Urréa Valencia, Valéria Ferreira Lopes, companheiros das minhas coletas.

Agradeço ao amigo Rodrigo Endres Ardissonne pelos bons momentos de discussões científicas, filosóficas e outras nem tanto, regadas com muitas risadas.

Ao professor Rafael Trevisan pelos ensinamentos e agradáveis conversas.

A Silvia pela amizade e companheirismo.

Agradeço aos demais professores do PPGFAP e colegas do mestrado, pelos ensinamentos e companheirismo.

Ao meu pai, por me incentivar a estudar sempre e mostrar que o conhecimento é a base do sucesso.

Em especial à minha família, por apoiar este sonho pessoal com compreensão, carinho e amor.

Agradeço ao Programa de Pós-Graduação em Biologia Vegetal (PPGFAP) da Universidade Federal de Santa Catarina, e a CAPES - Coordenação de Aperfeiçoamento de Pessoal de Nível Superior pelo auxílio financeiro sob a forma de bolsa e pela possibilidade de conhecer pesquisadores de diferentes instituições brasileiras e pela experiência adquirida em diferentes cursos.





## RESUMO

Políporos é um dos principais grupos de fungos decompositores de madeira, contudo, o conhecimento sobre sua diversidade ainda é escasso. Em Santa Catarina, a sinopse sobre a diversidade dos macromicetos apresenta 247 espécies de fungos, destas, 143 são políporoides, principalmente da região insular. Para a continuidade e o avanço do conhecimento sobre a diversidade de fungos políporoides que ocorrem na Mata Atlântica de Santa Catarina, foram estudados aspectos sobre a ecologia de comunidades fúngicas, particularmente de uma área pouco explorada, bem como um complexo taxonômico que ocorre na região. Primeiramente foi avaliada e caracterizada a comunidade de políporos em uma área de Floresta Ombrófila Densa do Parque Nacional da Serra do Itajaí, em termos de riqueza, abundância e funcionalidade. Entre os 152 espécimes coletados 58 espécies foram identificadas. *Polyporus dictyopus*, *Perenniporia martia* e *Fuscoporia walhbergii* foram as espécies dominantes. Com base nas frequências relativas nos diferentes tipos e substratos foram encontrados cinco grupos funcionais, dos quais dois formados por *Phylloporia spathulata* (em raízes vivas no solo) e *Phylloporia chrysa* (em lianas vivas) e os outros três constituídos por espécies saprofíticas, sendo cada grupo caracterizado pela presença de uma das espécies dominantes citadas. Finalmente, *Phellinusiptadeniae* é apresentada como um complexo taxonômico, em processo de especiação, a partir de evidências ecológicas (hospedeiro e distribuição geográfica) e filogenéticas.

**Palavras-chave:** Comunidade de Políporos Degradadores de Madeira; Grupos funcionais; Complexo Taxonômico *Phellinusiptadeniae*.



## ABSTRACT

Polypores is one of the most important groups of wood-decay fungi; however, its diversity knowledge is still scarce. In Santa Catarina state, a synopsis of macrofungi diversity presented 247 species, of which 143 are polypores, mainly from insular region. In order to continue and improve the knowledge about the diversity of polypores, which are occurring in the Atlantic Forest of Santa Catarina, ecological aspects of fungal communities were studied, in an unexplored area, as well as a complex taxonomy that occurs in the region. Firstly, was described the community of wood-decaying polypores in the Atlantic Rain Forest of Parque Nacional da Serra do Itajai, characterizing the community in terms of richness, abundance and functionality. Among 152 specimens collected 58 species were identified. *Polyporus dictyopus*, *Perenniporia martia* and *Fuscoporia wallbergii* were the dominant species. Based on the relative frequency in the different types of substrates where five functional groups were recognized. Two of them were formed by *Phylloporia spathulata* (on live roots in the ground) and *Phylloporia chrysa* (on live trunks); and three other functional groups, consisting of saprophytic species, each of them was characterized by the presence of one of the dominant species. Finally, *Phellinus iptadeniae* is presented as a taxonomic complex, in a speciation process, evidenced by the existence of two phylogenetic and ecological (host and geographic distribution) lineages.

**Keywords:** Wood-decay Polypore Communities; Functional Groups; *Phellinus iptadeniae* Complex.



## LISTA DE FIGURAS

### **Artigo 1 –**

Figure 1 - Map of collection area in the National Park of the Serra de Itajai

Figure 2 - Graph of rarefaction curve

Figure 3 - Graph of abundance/dominance curves

Figure 4 - Graph richness of species in different types of substrates.

Figure 5 - Graph of functional groups diameter only.

### **Artigo 2 –**

Figure 1 - Basidiomes of *Phellinus grenadensis* and *Phellinus piptadeniae*

Figure 2 - Skeletal hyphae of *Phellinus grenadensis* and *Phellinus piptadeniae*

Figure 3 - Phylogenetic tree



## LISTA DE TABELAS

### Artigo 1

Table 1 - Total relative frequency and number of polypores species in the substrate conditions and mean volume of the logs where they were found.

### Artigo 2

Table 1. Information of partial ITS rDNA sequences used in the phylogenetic analysis.

Table 2 - Morphological, host and geographical distribution data of *Phellinus grenadensis* (type specimen) and *Phellinus piptadeniae*.





## LISTA DE ABREVIATURAS E SIGLAS

- Alt. - Altitude.
- ARF - Atlantic Rain Forest.
- BE - do inglês broadly elipsoid, fortemente elipsoide.
- Caa - Caatinga.
- cm - centímetros.
- DB - Dead branche.
- DT - Dead trunk.
- Di - dimitic.
- et al. - do latim *et alii*, e outros, e colaboradores.
- Fig., Figs. - Figura, Figuras.
- i.e. - do latim *id est*, isto é.
- ibid. - do latim *ibidem*, no mesmo lugar.
- IKI - Reagente de Melzer.
- KOH - Hidróxido de Potássio, em solução a 3-5%.
- LT - Living trunk.
- m - metros.
- m.a.s.l. - do inglês meters above sea level, metros acima do nível do mar.
- mm - milímetros.
- Mono - monomitic.
- PE - Pernambuco.
- OG - On the ground.
- s. l. - do latim *sensu lato*, no sentido amplo.
- SC - Santa Catarina.
- SP - São Paulo.



## SUMÁRIO

INTRODUÇÃO GERAL .....	21
OBJETIVOS .....	23
LITERATURA.....	24
CAPÍTULO I. Community Structure and Functional Diversity of Polypores (Basidiomycota) in the Atlantic Rain Forest of Santa Catarina State, Brazil .....	27
INTRODUCTION.....	30
MATERIAL AND METHODS .....	31
Study area.....	31
Sampling methods .....	31
Data analysis .....	34
RESULTS .....	35
Species richness .....	35
Community structure: abundance and richness of species .....	36
Community structure: Functional groups.....	38
DISCUSSION .....	43
LITERATURE CITED .....	44
CAPÍTULO II. <i>Phellinus piptadeniae</i> complex: ecological, morphological and molecular evidences of a speciation process in Brazil .....	48
INTRODUCTION.....	51
MATERIAL AND METHODS .....	51
RESULTS AND DISCUSSION .....	55
LITERATURE CITED .....	63
CONSIDERAÇÕES FINAIS.....	66



## INTRODUÇÃO GERAL

As transformações impulsionadas pela expansão das fronteiras humanas, como crescimento das cidades, agricultura e pecuária, têm exercido forte pressão sobre o ambiente natural, com a destruição dos mais diferentes habitats, e conseqüentemente, levado a diminuição da diversidade biológica, muito ainda desconhecida. Esses fatos, entre outros, têm tornado a biodiversidade um dos assuntos mais relevantes das últimas décadas (Wilson 1994). Excluindo-se os insetos, os fungos constituem o mais numeroso grupo de seres vivos existentes, encontrados em praticamente todos os ecossistemas. Nesse contexto, estima-se que existam em torno de 1.5 a 5.1 milhões de espécies de fungos (Hawksworth 2001, Blackwell 2011), sendo que destas apenas cerca de 100 mil espécies foram descritas até o momento (Kirk *et al.* 2008).

O bioma Mata Atlântica vem sofrendo historicamente com a rápida degradação de ecossistemas e conseqüente eliminação de espécies. Este processo é resultado do crescimento da população humana sem limites e ocupação indevida de áreas que deveriam ser preservadas. Esta degradação deixou marcas profundas e atualmente o bioma é representado apenas por fragmentos remanescentes dos diferentes ecossistemas do bioma (Chapron 2010).

Em Santa Catarina, a sinopse sobre a diversidade dos macromicetos apresenta 247 espécies de fungos, sendo 33 de *Ascomycota* Whittaker e 212 de *Basidiomycota* R.T. Moore. Destes, 143 são poliporoides *s.l.* (*Hymenochaetales* Oberw., *Polyporales* Gäum. e *Trechisporales* K.H. Larss) e a grande maioria dos registros é proveniente da região insular (Loguercio-Leite *et al.* 2009). Considerando que há mais de 4.000 espécies de Angiospermas na Mata Atlântica de Santa Catarina (Forzza *et al.* 2010) e se utilizarmos a proporção fungos/plantas conservadora de 2:1 (Mueller *et al.* 2006), é possível estimar que o número de espécies de macromicetos possa ser de 8.000 para o Estado. Desta forma, as 143 espécies de poliporoides *s.l.*, principais agentes lignolíticos dentre os fungos degradadores de madeira, representariam pouco mais de 1% do total.

Em relação à diversidade funcional deste grupo de fungos pouco ainda se conhece sobre seus aspectos ecológicos. As espécies de políporos apresentam afinidades distintas por

diferentes substratos, que variam desde ramos de pequeno diâmetro até troncos grandes de árvores vivas. Tem sido proposto que é possível reconhecer grupos funcionais de fungos lignolíticos baseado nas relações de preferências por tipos de substrato em particular. Estes grupos apresentam capacidade de degradação diferenciada e assim estariam cumprindo diferentes funções no ecossistema (Urcelay & Robledo 2004). O conceito de grupos funcionais sustenta a separação das espécies de fungos poliporoides em categorias que estão diretamente relacionadas com o tipo de substrato (troncos e ramos/galhos de árvores vivas ou mortas, em pé ou ainda ao solo e de diferentes diâmetros), bem como os diferentes estágios de decomposição. Estudos recentes, realizados na Argentina, evidenciam que os fungos polipóridos podem ser classificados em grupos funcionais (Urcelay & Robledo 2004; Robledo 2009).

No Brasil, a Mata Atlântica é um mosaico diversificado, apresentando estruturas e composições florísticas distintas, devido às diferenças de solo, topografia e clima, características disponíveis na ampla área de ocorrência desse bioma no Brasil (IBAMA, 2008). O bioma tem uma diversidade de espécies de plantas calculada em 20.000 espécies (Capobianco 2001; Mittermeier, 1999), dos quais 55% são árvores, proporcionando um grande número de nichos onde as variáveis como a umidade, temperatura, volume e diâmetro dos troncos e ramos, estágios de decomposição, estrutura físico-química das diferentes de árvores mortas (Rayner & Todd, 1979) podem abrigar diferentes grupos funcionais e diferentes estruturas nas comunidades de fungos lignolíticos.

Outro aspecto que subestima a diversidade de políporos é a ocorrência de espécies crípticas, ou seja, complexos taxonômicos que podem ou não apresentar diferentes espécies com morfologia similar. Para a América do Sul, Decock *et al.* (2007) registraram a existência destas espécies crípticas em *Hymenochaetales* poroides, táxons que não apresentam diferenças morfológicas perceptíveis. Para a resolução destes complexos são necessárias análises complementares à morfológica, ou seja, uma abordagem mais integrativa utilizando também caracteres ecológicos, biogeográficos, moleculares e de biologia reprodutiva. Na ilha de Santa Catarina, os basidiomas himenquetoides (pileados e poroides) coletados em *Piptadenia gonoacantha* (Mart.) J.F.Macbr., foram tradicionalmente determinados como *Phellinus grenadensis* (Murrill) Ryvardeen (Drechsler-Santos & Loguercio-Leite 2006). Esta espécie foi descrita para a ilha caribenha de Grenada, que de acordo com Ryvardeen (2004) é amplamente distribuída

pelos trópicos e em hospedeiros de diversos gêneros. *Piptadenia gonoacantha*, é uma planta popularmente chamada de “pau-jacaré” com ocorrência em todo o bioma Mata Atlântica. No estado de São Paulo, em 1950, o micólogo brasileiro A.R. Teixeira descreveu *Phellinus iptadeniae* Teixeira como uma espécie encontrada unicamente em “pau-jacaré”. Por sua vez, *Phellinus grenadensis* da ilha de SC apresenta morfologia muito similar ao material tipo de *P. iptadeniae*. Em suma, é possível que os materiais da ilha de SC determinados como *Phellinus grenadensis* correspondam à espécie *Phellinus iptadeniae*, que de acordo com Drechsler-Santos *et al.* (2010) ocorre também no Bioma Caatinga com níveis de especialização (*host-recurrent*) em espécies de *Piptadenia*.

Diante do apresentado, faz-se necessário avançar como o conhecimento sobre a diversidade de fungos políporos que ocorrem na Mata Atlântica de Santa Catarina, estudando especificamente aspectos sobre a ecologia das comunidades fúngicas, particularmente, em áreas pouco exploradas, mas também sobre a taxonomia de complexos de espécies existentes.

## OBJETIVOS

Com o objetivo principal de investigar aspectos ecológicos e taxonômicos dos fungos poliporoides (Basidiomycota) da Mata Atlântica de Santa Catarina, especificamente pretende-se: (i) Avaliar e caracterizar a comunidade de políporos em termos de riqueza, abundância e funcionalidade de fungos políporos em uma área de Floresta Ombrófila Densa do Parque Nacional da Serra do Itajaí; (ii) Reavaliar taxonomicamente *Phellinus grenadensis* da ilha de Santa Catarina com base na morfologia, ecologia e análises moleculares.

## LITERATURA

BLACKWELL, Meredith. The fungi: 1, 2, 3 ... 5.1 million species? *American Journal of Botany* 98(3): 426–438. 2011.

CAPOBIANCO João Paulo Ribeiro, et al. *Biodiversidade na Amazônia Brasileira*. Estação Liberdade & Instituto Sócio-Ambiental, São Paulo. 540p. 2001.

CHAPRON, Guillaume. 2010. Biodiversity: from conservation science to action. *SOAPBOX SCIENCE*  
<<http://blogs.nature.com/soapboxscience/2010/09/10/>>. acesso em 01/03/2013.

DECOCK Cony, et al. *Fomitiporia punctata* (Basidiomycota, Hymenochaetales) and its presumed taxonomic synonyms in America: taxonomy and phylogeny of some species from tropical/subtropical areas. *Mycologia* 99(5):733-52. 2007.

DRECHSLER-SANTOS, Elisandro Ricardo, LOGUERCIO-LEITE, Clarice. Basidiomycetos SC – Brasil. *Guia Eletrônico de Identificação. Version BASC – 1.0*. 2006. <http://www.ccb.ufsc.br/bot/micologia/>  
Acesso:em 20/06/2011.

DRECHSLER-SANTOS, Elisandro Ricardo et al. Ecological aspects of Hymenochaetaceae in an area of Caatinga (semi-arid) in Northeast Brazil. *Fungal Diversity* 42: 71-78. 2010.

FORZZA, Rafaela Campostrini, et al. *Angiospermas in Lista de Espécies da Flora do Brasil*. Jardim Botânico do Rio de Janeiro. 2010. (<http://floradobrasil.jbrj.gov.br/2010/FB000032>) acesso em 01/03/2012.

HAWKSWORTH, David Leslie. The magnitude of fungal diversity: the 1.5 million species estimate revisited. *Mycological Research* 105(12): 1422-1432. 2001.

KIRK, Paul Michael, et al. *Ainsworth & Bisby's Dictionary of the Fungi*, 10. ed. Wallingford, Oxon.: CAB international, 771 p. 2008.



LOGUERCIO-LEITE, Clarice, et al. Résumé of macromycetes from Santa Catarina State, Southern Brazil. *Ínsula* (Florianópolis): 38: 1-14. 2009.

MITTERMEIER, Russel Albert et al. *Hotspots: earts biologicaly richest and most endangered terrestrial ecoregions*. Mexico City:CEMEX. 430p. 1999.

MUELLER, Gregory Michael et al. Global diversity and distribution of macrofungi. *Biodiversity and Conservation* 16: 37-48. 2006.

RAYNER, ADM, TODD, NK. Population and community structure and dynamics of fungi in decaying wood. *Adv. Bot. Res.* 7, 333–420. 1979.

RYVARDEN, Lief. *Neotropical Polypores Part 1*. Oslo, Fungiflora. 2004.

URCELAY, Carlos; ROBLEDO, Gerardo. Community structure of polypores (Basidiomycota) in Andean alder wood in Argentina: Functional groups among wood-decay fungi? *Austral Ecology*, p. 471-476. 2004.

URCELAY, Carlos; ROBLEDO, Gerardo. Positive relationship between wood size and basidiocarp production of polypore fungi in *Alnus acuminata* forest. *Fungal Ecology*, v. 2, n. 3, p. 135-139. 2009.

WILSON, Edward Osborne. *Diversidade da Vida*. São Paulo. Cia da Letras, 447p. 1994.



CAPÍTULO I. Community Structure and Functional Diversity of  
Polypores (Basidiomycota) in the Atlantic Rain Forest of Santa Catarina  
State, Brazil

**Community structure and Functional diversity of Polypores (Basidiomycota) in the Atlantic Rain Forest of Santa Catarina State, Brazil.**

*Marco A. Borba-Silva<sup>1\*</sup>; Gerardo L. Robledo<sup>2</sup> & E. Ricardo Drechsler-Santos<sup>1</sup>*

*1. Universidade Federal de Santa Catarina, Departamento de Botânica, Campus Universitário, Trindade, Florianópolis, SC, CEP: 88040-900.*

*2. Laboratorio de Micología, Instituto Multidisciplinario de Biología Vegetal, CONICET - Universidad Nacional de Córdoba, CC 495, CP 5000 Córdoba, Argentina*

*\*Correspondence to: [marcoborba64@terra.com.br](mailto:marcoborba64@terra.com.br)*

## ABSTRACT

Ecological studies have suggested that different groups of polyporoid species, acting as parasites and/or saprophytes, degrade different types of woody substrates. These functional groups have different decay capabilities and hence different roles in the ecosystems. The aim of this study was to describe the community (species composition and functional groups present) of wood-decaying polypores in the Atlantic Rain Forest of Parque Nacional da Serra do Itajaí, in the Santa Catarina state. The polypores specimens and substrate data were collected in two plots (100x50m). Among 152 specimens collected 58 species were identified. Three dominance groups were identified. A first group of three dominant species (*Polyporus dictyopus*, *Perenniporia martia* and *Fuscoporia walhbergii*); a second group of five subordinate species and a third group of 50 rare species. The species were ordered using a cluster correspondence analysis based on relative frequency in the different types of substrates and the mean size of the substrate they fruit. Five functional groups were recognized: two of them are formed by *Phylloporia spathulata* (on live roots in the ground) and *Phylloporia chrysitia* (on live trunks); and three other functional groups, consisting of saprophytic species, each of them was characterized by the presence of one of the three dominant species, subordinate and rare ones. The importance of ecological studies addressing community structure is discussed.

## INTRODUCTION

Polypores constitute one of the main groups of wood-decay fungi. Although their importance in the ecology of forests, the diversity of wood-decay fungi of the Atlantic Rain Forest (ARF) has been approached by several authors (see for instance Loguercio-Leite *et al.* 2002; Ryvarden & de Meijer 2002; Drechsler-Santos *et al.* 2008). Recently Baltazar & Gibertoni (2009) presented the most extensive literature survey of 733 apylophoraceous species, of which more than 50% are polypores.

Regarding the functional diversity of this group of fungi, little is known about its ecological aspects. Ecological studies have suggested that different groups of polyporoid species, acting as parasites and/or saprophytes, degrade different types of woody substrates. These functional groups have distinct capacity degradation and perform different roles in the ecosystems. Among the wood-decay fungi *Polyporaceae s.l.*, as well as some *Corticaceae s.l.* and *Hymenochaetaceae*, have been recommended as well suited to study the key role of woody debris in old-growth forests (Bader *et al.* 1995).

In this context, Urcelay & Robledo (2004) have proposed that it is possible to recognize functional groups of polypores based on the preferences of the species for particular types of substrate. The authors recognized groups of species that are directly related to different type of substrates (trunks and branches/twigs of trees living or dead, standing or on the ground and of different volumes) as well as different decay stages. This functional group proposal is based in a study conducted in monospecific *Alnus acuminata* H.B.K (Betulaceae) forests of Northwest Argentinean Yungas.

The polypores functional groups, proposed by Urcelay & Robledo (2004), should be tested in a highly diverse ecosystem, as Atlantic Rain Forest of Brazil. This biome is a diversified mosaic, showing structures and floristical compositions distinguished, due to the differences in soil, topography and climate characteristics available in the broad area of occurrence in Brazil (IBAMA, 2008). The plant species diversity estimated of Atlantic Rain Forest is around 20.000, of which 55% are trees (Capobianco 2001; Mittermeier, 1999). This high diversity provides a large number of niches with different variables (humidity, temperature, volume and diameter of trunks and branches, stage of decomposition of wood, and physical and chemical structure of dead trees) that may influence in the functional diversity and community structure of wood-decay fungi (Rayner & Todd 1979).

The aim of this study is to describe and characterize the community of wood-decay polypores of the Parque Nacional da Serra do Itajaí (PARNA-SJ), in the Atlantic Rain Forest of Santa Catarina state, Brazil, in terms of richness, abundance and functionality.

## MATERIAL AND METHODS

### Study area

The Parque Nacional da Serra do Itajaí (PARNA-SI) is located in the Itajaí Valley, in Santa Catarina state (Figure 1A), covering an area of 57.374 ha and altitudes from 80 to 1039 m.a.s.l. The climate is temperate humid hot summer (Cfa), no deficiency of rain in any season with evapotranspiration with potential mega thermal (GAPLAN, 1986). The annual mean temperature ranges between 17 ° C and 22 ° C, having an annual rainfall ranging between 1600 and 1800 mm, distributed between 120 and 140 days of rain during the year, with an average relative humidity between 75 and 85% (Klein, 1979). This conservation area comprises a significant portion of the remnants of the Atlantic Rain Forest biome, with patches of primary forests and others in advanced stage of regeneration (source: SOS Mata Atlântica and INPE Instituto Socioambiental). The study was conducted in the Parque Municipal Natural Nascentes do Garcia, Blumenau municipality, where two study areas of 5000 m<sup>2</sup> (100 x 50 m) were plotted (Figure 1B). The vegetation in the study area is classified as Dense Ombrophilous Forest (GAPLAN 1986).

### Sampling methods

Each plots (Figure 1B) was sampled four times each one (field expeditions: 5-6 November 2011, 22-23 February 2012 , 11-12 June 2012 and 12-13 September 2012) in order to cover the seasons of one year. Figure 1 shows the area and the position of the two plots.


All basidiomes of polypores species were recorded and when there were many basidiomes of the same species in the same substrate only one occurrence was recorded.. Most of substrate sampled was identified by photos in order to avoid recording the same specimens during different field expeditions. The following substrate data were recorded for each fungal occurrence: diameter and length (volume of the substrate was then calculated) and the condition [living trunk (LT); dead trunk (DT): standing or fallen logs and stumps; living branch (LB); dead

branch (DB); and on the ground (OG)]. To consider the trunks established the need for visualization of stumps near the trunks.

According to Urcelay & Robledo (2004), in the present study it was assumed that there was a gradient in the diameter and volume of the logs and a gradient in the level of decay, ranging from the dead trunks trunks living in the soil, representing a decay succession. In the present study we use this classification to decay wood: initial (the bark rot and decay on wood up to 3cm), moderate (the cortical portion with wood rot more than 3 cm, with the center of the timber a hard consistency) and advanced (substrate can be removed easily, all be rotten wood).



**Figure 1.** Study area: (A) PARNA Serra do Itajaí in the Santa Catarina state, with delimitation of study area (the square represents the Parque Natural Nascentes do Garcia extended in B); (B) Parque Natural Nascentes do Garcia showing the position of plots.



## Data analysis

Estimated richness was calculated with ACE and CHAO 1 estimators with one individual-based frequency sample (all polypore species frequency records across all sampling plots and dates) with the software EstimateS (Version 9.0, [viceroy.eeb.uconn.edu/estimates](http://viceroy.eeb.uconn.edu/estimates/)).

The relative frequency of each polypore species in each substrate condition was calculated as:

$$Fr_{xi} = \frac{n_{xi}}{A_x}$$

where  $n_{xi}$  = number of occurrences of each polypore species  $x$  in each substrate condition  $i$  (*viz.*, LT, DT, LB, DB, and OG) and  $A_x$  = abundance of polypore species  $x$  (across all sampling plots and dates). The species richness in each substrate condition was also calculated.

Cluster analyses were performed to classify polypores species into groups as recommended by Goodall (1980). This method is based on distances between species and combines the species computed from their relative frequency on different substrate conditions ( $Fr_{xi}$ ) and the average volume of the substrates on which they occurred (Table 1). The mean Euclidean distance was used as a measure of similarity between species and clustering was based on the average linkage method using the Infostat statistical package (Di Rienzo *et al.* 2001). Species with two or more occurrences were included in the analysis (Urcelay & Robledo 2004).

The sampling effort was tested using the complementarity (dissimilarity), an empirical measure of the degree of species uniqueness between different samples. This empirical test for complementarity was calculated as:

$$C_{jk} = \frac{U_{jk}}{S_{jk}}$$


where  $C_{jk}$  = the complementarity between samples (or plots)  $j$  and  $k$ ,  $U_{jk}$  = the total number of taxa encountered at only one plot, and  $S_{jk}$  is the total richness of both plots combined (Colwell & Coddington 1994). Sampling effort for species richness is most efficient when complementarity among samples ( $C_{jk}$ ) is about 0.5 (Lodge *et al.* 2004).

## RESULTS

### Species richness

In the present study 152 records were done, from which 94 % (143 records) were identified at species level representing 58 polypore species. Eleven specimens (6%) could not be identified at species level, however they represent 11 distinct species among the polypore community (i.e. *Dichomitus sp.*, *Fomitiporella sp.*, *Fuscoporia sp.*, *Junghuhnia sp.*, *Oxyporus sp.*, *Phellinus sp.1* and *sp.2*, *Polyporus sp.*, *Rigidoporus sp.*, *Tyromyces sp.1* and *sp. 2*), and therefore have been included in the study. Species richness observed (58 species) represent 51% of the estimated richness to ACE=114.1 and 47% to CHAO1=122.2 (Fig. 2). Complementarity (dissimilarity) was  $C_{jk} = 0.7$ , showing that the sampling for richness was efficient.


**Figure 2.** Observed S(est) and Estimated richness by CHAO1 and ACE estimators. Estimators show different values.



### Community structure: abundance and richness of species


Community structure based on species abundance shows (Figure 3, Table 1) that *Polyporus dictyopus*, *Perenniporia martia*, and *Fuscoporia walhbergii* were the dominant species showing the highest abundance values (15, 13 and 12 respectively). They represent almost 26% of the total occurrences (40 out of 152) but only 5.4% of the species richness. In the other extreme, a group of rare species, mainly with one, two or three records (only two with four records) represent around 50% of the total of occurrences (76/152); however they cover the 86% of species richness. With intermediate abundance values of six to nine records, a group of five subordinate species represent 24% of the total occurrences (38/152) and 8.6% of species richness.

**Figure 3.** Abundance of species of the PARNA Serra do Itajaí: **D.** dominant species; **S.** subordinate species and **R.** rare species.



Abundance and species richness were increasing according the advanced decay stage of the substrate. Among 152 specimens collected, 6 (4 species) were recorded on LT, 68 (22 species) on DT, 71 (40 species) on DB and 7 (5 species) on the ground (OG) (Figure 4, Table 1).

**Figure 4.** Richness and relative abundance of polypore species of the PARNA Serra do Itajaí within different substrate types.



#### Community structure: Functional groups


Based on the relative frequency of each species in the different log conditions and the mean diameter and volume of the log on which the basidiomas were found, the cluster analysis produced four main groups of polypore species when the dendrogram was cut at a distance of 2,30 (Figure 5). *Phylloporia spathulata* (with relative frequency of 1.00 on OG) and *Phylloporia chrysitata* (with relative frequency of 1.00 in LT) formed two functional groups, Group 1 and 2 respectively, themselves.

Group 3 is formed by *Ganoderma australe*, *Phellinus detonsus* and *Fuscoporia wahlbergii*. These species present the capability to

decay and basidioma emerge in standing living trunks, but also can continue growing as saprobes and the basidioma emerge on dead trunks too.

Groups 4 and 5 are constituted by saprobe species decaying dead wood, and they are differentiated by the relative frequency and the average size of the substrates. Group 4 is formed by seven species that grow “preferably” on dead trunks of large size, the highest volume values observed are in this group. Group 5 comprise 18 species that grow exclusively on dead branches of small volume. Within Group 5, two smaller subgroups could be distinguished. *Rigidoporus lineatus*, *Steccherinum reniforme*, *Trichaptum sector* and *Polyporus dictyopus* form a subgroup characterized by grow indifferently on DT and DB of relatively low volume. The other subgroup is composed by species that grows exclusively in DB with the lowest volume values. Regarding groups 3 to 5, each group is composed by 1 dominant species, at least one subordinate and several rare species (only one in group 3).

**Figure 5.** Cluster analysis of different polypores species of the PARNA Serra do Itajaí based on relative frequency of the species in each substrate condition and the mean volume of the substrates where they were found. Arrows indicates dominant species.





**Table 1.** Total relative frequency and number of polypores species in the substrate conditions and mean volume of the logs where they were found.

Species	Relative Frequency						$\bar{X}Vol$ (m <sup>3</sup> )
	A	OG	LB	LT	DB	DT	
<i>Polyporus dictyopus</i>	15	0,00	0,00	0,00	0,67	0,33	0,175
<i>Perenniporia martia</i>	13	0,00	0,00	0,00	0,00	1,00	1,567
<i>Fuscoporia wahlbergii</i>	12	0,00	0,00	0,08	0,00	0,92	0,326
<i>Fomitella supina</i>	9	0,00	0,00	0,00	0,33	0,67	0,744
<i>Fuscoporia gilva</i>	9	0,00	0,00	0,00	0,44	0,56	0,935
<i>Rigidoporus microporus</i>	7	0,00	0,00	0,00	0,29	0,71	1,094
<i>Ganoderma australae</i>	6	0,00	0,00	0,33	0,00	0,67	0,056
<i>Steccherinum reniforme</i>	6	0,00	0,00	0,00	0,67	0,33	0,247
<i>Corioloopsis caperata</i>	4	0,00	0,00	0,00	0,25	0,75	1,168
<i>Rigidoporus lineatus</i>	4	0,00	0,00	0,00	0,50	0,50	0,022
<i>Pachykytospora alabamae</i>	3	0,00	0,00	0,00	1,00	0,00	0,001
<i>Phylloporia spathulata</i>	3	1,00	0,00	0,00	0,00	0,00	0,000
<i>Polyporus tenuiculus</i>	3	0,00	0,00	0,00	1,00	0,00	0,002
<i>Pycnoporus sanguineus</i>	3	0,00	0,00	0,00	1,00	0,00	0,025
<i>Trichaptum sector</i>	3	0,00	0,00	0,00	0,67	0,33	0,029
<i>Dichomitus cylindrosporus</i>	2	0,00	0,00	0,00	1,00	0,00	0,002
<i>Junghuhnia polycistidifera</i>	2	0,00	0,00	0,00	0,50	0,50	0,570
<i>Phellinus detonsus</i>	2	0,00	0,00	0,50	0,00	0,50	0,173
<i>Phylloporia chrysitae</i>	2	0,00	0,00	1,00	0,00	0,00	0,001
<i>Polyporus guianensis</i>	2	0,00	0,00	0,00	1,00	0,00	0,001
<i>Polyporus sp.</i>	2	0,00	0,00	0,00	1,00	0,00	0,001
<i>Polyporus tricholoma</i>	2	0,00	0,00	0,00	1,00	0,00	0,001
<i>Rigidoporus sp.</i>	2	0,00	0,00	0,00	0,50	0,50	1,768
<i>Trametes villosa</i>	2	0,00	0,00	0,00	1,00	0,00	0,001
<i>Abortiporus sp.</i>	1	0,00	0,00	0,00	1,00	0,00	0,001
<i>Amauroderma schomburgkii</i>	1	1,00	0,00	0,00	0,00	0,00	0,000
<i>Amauroderma sp.</i>	1	1,00	0,00	0,00	0,00	0,00	0,000
<i>Amauroderma sprucei</i>	1	1,00	0,00	0,00	0,00	0,00	0,000
<i>Antrodiella sp.</i>	1	0,00	0,00	0,00	1,00	0,00	0,001
<i>Ceriporia mellea</i>	1	0,00	0,00	0,00	1,00	0,00	0,001

<i>Ceriporiopsis balaenae</i>	1	0,00	0,00	0,00	1,00	0,00	0,016
<i>Ceriporiopsis latemarginata</i>	1	0,00	0,00	0,00	1,00	0,00	0,006
<i>Coltricia barbata</i>	1	1,00	0,00	0,00	0,00	0,00	0,000
<i>Corioloopsis rigida</i>	1	0,00	0,00	0,00	1,00	0,00	0,011
<i>Cystidiodontia laminifera</i>	1	0,00	0,00	0,00	0,00	1,00	0,377
<i>Dichomitrus sp.</i>	1	0,00	0,00	0,00	1,00	0,00	0,003
<i>Diplomitoporus venezuelinicus</i>	1	0,00	0,00	0,00	0,00	1,00	1,256
<i>Flabellophora parva</i>	1	0,00	0,00	0,00	1,00	0,00	0,000
<i>Fomitiporella sp.</i>	1	0,00	0,00	0,00	1,00	0,00	0,004
<i>Fuscoporia puntactiformes</i>	1	0,00	0,00	0,00	1,00	0,00	0,001
<i>Fuscoporia sp.</i>	1	0,00	0,00	0,00	0,00	1,00	0,791
<i>Gramothele sp.</i>	1	0,00	0,00	0,00	1,00	0,00	0,001
<i>Henningsia brasiliensis</i>	1	0,00	0,00	0,00	0,00	1,00	0,006
<i>Hexagonia hydnooides</i>	1	0,00	0,00	0,00	1,00	0,00	0,003
<i>Junghuhnia sp.</i>	1	0,00	0,00	0,00	1,00	0,00	0,001
<i>Lentinus crinitus</i>	1	0,00	0,00	0,00	1,00	0,00	0,001
<i>Oxiporus pellicula</i>	1	0,00	0,00	0,00	1,00	0,00	0,001
<i>Oxiporus sp.</i>	1	0,00	0,00	0,00	1,00	0,00	0,045
<i>Phellinus melleoporus</i>	1	0,00	0,00	0,00	0,00	1,00	1,360
<i>Phellinus sp.1</i>	1	0,00	0,00	0,00	0,00	1,00	0,109
<i>Phellinus sp.2</i>	1	0,00	0,00	0,00	0,00	1,00	0,035
<i>Phellinus umbrinellus</i>	1	0,00	0,00	0,00	0,00	1,00	0,814
<i>Polyporus arcularius</i>	1	0,00	0,00	0,00	1,00	0,00	0,026
<i>Polyporus gramocephalus</i>	1	0,00	0,00	0,00	1,00	0,00	0,014
<i>Rigidoporus undatus</i>	1	0,00	0,00	0,00	1,00	0,00	0,008
<i>Tyromyces sp.1</i>	1	0,00	0,00	0,00	1,00	0,00	0,001
<i>Tyromyces sp.2</i>	1	0,00	0,00	0,00	1,00	0,00	0,004
<i>Wrightoporia bracei</i>	1	0,00	0,00	0,00	1,00	0,00	0,001
<b>Species richness</b>		5	0	4	39	23	

## DISCUSSION

This work was based on the occurrence of species through presence of their basidiomes. The results and conclusions were underestimated for a part of the community, because they did not have basidiomes on substrate and were not considered. We conducted the sampling through four seasons in order to consider the seasonal variability and to include as many species as possible. Work with presence-absence of basidiomes is a methodology accepted to do ecological works (Urcelay & Robledo 2004 and references there in).

Species richness and abundance increase as increases the decay stage as here is assumed. This is in accordance with previous results (Urcelay & Robledo 2004) and the theoretical frame of decay succession (Rayner & Todd 1979).

Regarding functional structure of the community, our results agree with those of Urcelay & Robledo (2004). We could identify two additional functional groups (Group 1 and 2), both constituted by *Phylloporia* species. *Phylloporia* species are parasites of roots, living branches and living stems of creepers (Decock *et al.* 2013). This genus is not present in the *Alnus acumniata* montane forests where polypore functional groups were originally proposed (Robledo *et al.* 2003, Urcelay & Robledo 2004). The groups 3-5 identify in this study agree with those proposed by Urcelay & Robledo (2004), however the species composition is different. This support the idea that all woody substrates types and sizes are important for fungal diversity (Heilmann-Clausen & Christensen 2004).

Species richness observed (58) is near 50% of the estimated (ACE=114.1 and CHAO1=122.2). This represent less than 16% and 31% to ACE and 33% to CHAO1 of ca. 365 polypores species reported by Baltazar & Gibertoni (2009) for the ARF biome. Sampling effort was enough as suggested by complementarity index (Lodge *et al.* 2004); abundance and richness values will increase certainly with more sampling efforts (additional plots), however we believe that dominance and functional structure will not change substantially. From a practical point of view the sample design of this study was enough to have a preliminary overview of the community structure.

Ecological studies describing the community structure of species, as here presented, are extremely important, particular actual conservation issues that threaten the remains of forests. Particular substrates and/or key functional species could be identified, and then could be considered when conservation decisions are taken.

## LITERATURE CITED

Bader P, Jansson S, Jonsson BG, 1995. Wood-inhabiting fungi and substratum decline selectively logged boreal spruce forests. *Biological Conservation* 72:355-362.

Baltazar JM, Gibertoni TB, 2009. A checklist of the aphylloroid fungi (Basidiomycota) recorded from the Brazilian Atlantic Forest. *Mycotaxon* 109: 439-442.

Capobianco JPR (organ.), 2001. Dossie Mata Atlântica 2001. Projeto Monitoramento Participativo da Mata Atlantica. Instituto Socioambiental (ISA), São Paulo. 15pp. 1 map.

Colwell RK, Coddington JA. 1994. Estimating terrestrial biodiversity through extrapolation. *Philosophical Transactions of the Royal Society London, B* 345:101–118

Colwell RK, 2013. EstimateS: Statistical estimation of species richness and shared species from samples. Version 9.0. Persistent URL <[purl.oclc.org/estimates](http://purl.oclc.org/estimates)>

Decock C, Amalfi M, Robledo G, Castillo G. (2013). *Phylloporia nouraguensis*, an undescribed species on Myrtaceae from French Guiana. *Cryptogamie Mycologie* 34 (1): 15-27.

Di Rienzo JA, Casanoves F, Balzarini MG, Gonzalez L, Tablada M, Robledo CW, 2012. InfoStat versión. InfoStat Group, Facultad de Ciencias Agropecuarias, Universidad Nacional de Córdoba, Argentina. URL <http://www.infostat.com.ar>

Drechsler-Santos ER, Wartchow F, Baseia IG, Gibertoni TB, Cavalcanti MAQ, 2008. Revision of the Herbarium URM I. Agaricomycetes from the semi-arid region of Brazil. *Mycotaxon* 104:9–18

Forzza RC, Leitman P, Walter BMT, Costa A, Pirani Jr, Morim MP, Queiroz LP, Martinelli G, Peixoto AL, Coelho MAN, Stehmann Jr, Baumgratz JFA, Lohmann LG, Hopkins M, 2010. Angiospermas in

Lista de Espécies da Flora do Brasil. Jardim Botânico do Rio de Janeiro. (<http://floradobrasil.jbrj.gov.br/2010/FB000032>).

Fundação SOS Mata Atlântica & Instituto Nacional De Pesquisas Espaciais, Instituto Socio-Ambiental. 2000. Atlas da evolução dos Remanescentes Florestais e Ecossistemas Associados no Domínio da Mata Atlântica no Período 1995-2000. São Paulo – SP.

GAPLAN. 1986. Atlas de Santa Catarina. Aerofoto Cruzeiro, Florianópolis.

Gilbertson RL, Ryvarde L, 1987. North American Polypores, Vol. 1-2. Fungiflora, Oslo.

Gilbertson RL, 1980. Wood-rotting fungi of North America. Mycologia 72, 1–49.

Goodall DW, 1980. Numerical classification. In: Classification of Plant Communities (ed. R. H. Whittaker) pp. 247–86. Kluwer Academic Publishers Group, The Hague.

Heilmann-Clausen J, Christensen M, 2004. Does size matter?: On the importance of various dead wood fractions for fungal diversity in Danish beech forests. Forest Ecology and Management 201 (1), 105-117.

Lindblad I, 1998. Wood-inhabiting fungi on fallen logs of Norway spruce: relations to forest management and substrate quality. Nord. J. Bot. 18, 243–55.

Lodge DJ, Ammirati JF, O'dell TE, Mueller GM, Huhndorf SM, Wang GJ, Stokland JN, Schmit JP, Ryvarde L, Leacock PR, Mata M, Umaña L, Wu Q (florence), Czederpiltz DL, 2004. Terrestrial And Lignicolous Macrofungi. In: Biodiversity of Fungi: Inventory and Monitoring Methods, Elsevier Academic Press. 777p.

Loguercio-Leite C, Ryvarde L, Groposo C, 2002. Studies in Neotropical polypores 16. *Rubroporus carneoporis* genus & species nova. Mycotaxon 83: 223-227.

Loguercio-Leite C, Campos-Santana M DE, Gerlach A, Gutjahr M, Trierveiler-Pereira L, Drechsler-Santos ER, Baltazar JM, 2009. Résumé

of macromycetes from Santa Catarina State, Southern Brazil. *Ínsula (Florianópolis)*: 38: 1-14.

Mittermeier RA, Myers N, Mittermeier CG, Gil PR, 1999. Hotspots: earth's biologically richest and most endangered terrestrial ecoregions. CEMEX & Conservation International. 431pp.

Mueller GM, Schmit JP, Leacock PR, Buyck B, Cifuentes J, Desjardin DE, Halling RE, Hjortstam K, Iturriaga T, Larsson KH, Lodge DJ, May TW, Minter D, Rajchenberg M, Redhead SA, Ryvar den L, Trappe JM, Watling R, Wu Q, 2007. Global diversity and distribution of macrofungi. *Biodiversity and Conservation* 16: 37-48.

Niemelä T, Renvall P, Pentillä R, 1995. Interactions of fungi at late stages of wood decomposition. *Ann. Bot. Fennici* 32, 141–52.

Oberwinkler F, 1994. Evolution of functional groups in Basidiomycetes (Fungi). In: *Biodiversity and Ecosystem Function* (eds E.-D. Schulze & H. A. Mooney) pp. 143–63. Springer-Verlag, Berlin.

Rayner ADM, Todd NK, 1979. Population and community structure and dynamics of fungi in decaying wood. *Adv. Bot. Res.* 7, 333–420.

Robledo GL, Urcelay C, Rajchenberg M, Domínguez L, 2003. Políporos (Aphylophorales, Basidiomycota) parásitos y saprófitos de *Alnus acuminata* en el noroeste argentino. *Boletín de la Sociedad Argetnina de Botánica* 38 (3-4): 207-224.

Ryvar den L, 2004. Neotropical polypores. Part 1: introduction, ganodermataceae & hymenochaetaceae. *Synopsis Fungorum* 19. Fungiflora, Oslo

Ryvar den L, Meijer AAR, 2002. Studies in neotropical polypores 14. New species from the state of Paraná, Brazil. *Synopsis Fungorum* 15: 34-69.

Urcelay C, Robledo G, 2004. Community structure of polypores (Basidiomycota) in Andean Alder wood in Argentina: Functional groups among wood-decay fungi? *Austral Ecology* 29: 471-476.



CAPÍTULO II. *Phellinus piptadeniae* complex: ecological, morphological and molecular evidences of a speciation process in Brazil



*Phellinus iptadeniae* complex

***Phellinus iptadeniae* complex: ecological, morphological and molecular evidences of a speciation process in Brazil**

M.A. Borba-Silva<sup>1\*</sup>, G.L. Robledo<sup>2</sup>, A. Góes-Neto<sup>3</sup> & E.R. Drechsler-Santos<sup>1</sup>  
 Universidade Federal de Santa Catarina, Departamento de Botânica, Campus  
 Universitário, Trindade, Florianópolis, SC, CEP: 88040-900.

\* Author for correspondence: marcoborba64@terra.com.br

M.A. Borba-Silva<sup>1\*</sup>, G.L. Robledo<sup>2</sup>, A. Góes-Neto<sup>3</sup> & E.R. Drechsler-Santos<sup>1</sup>  
 (Universidade Federal de Santa Catarina, Departamento de Botânica, Campus  
 Universitário, Trindade, Florianópolis, SC, CEP: 88040-900). *Phellinus  
 iptadeniae* complex: ecological, morphological and molecular evidences of a  
 speciation process in course in Brazil. J. Torrey The occurrence of cryptic  
 species has been documented among the Neotropical polypores. Species  
 complex, new species and speciation process were recently proposed using an  
 integrative approach (host and geographic distribution, morphological,  
 compatibility tests and molecular analysis). *Phellinus iptadeniae* was  
 described by A. Teixeira as specific on *Piptadenia gonoacantha* from Atlantic  
 forest of São Paulo State. The distribution of this species was expanded recently  
 to the Caatinga biome, where was characterized as recurrent on *Piptadenia spp.*  
 In the Atlantic Rain Forest of Santa Catarina state, specimens collected on  
*Piptadenia gonoacantha* have been traditionally named as *Phellinus  
 grenadensis*. A comparative morphological detailed study reveal that these  
 specimens correspond to *P. iptadeniae*, a fact also supported by molecular  
 analysis. Our results support that *Phellinus iptadeniae* is a taxonomic complex  
 of species that is undergoing a speciation process evidenced by the existence of  
 two phylogenetic and ecological (host and geographic distribution) lineages.

Key words –Taxonomic complex, *Hymenochaetaceae*, ITS analyses, Host  
 specificity.

The authors thank Dr. Carlos Urcelay (IMBIV, Argentina), Dr. Matheus A.  
 Reck (UFRGS) and Maria Alice Neves (PPGFAP/UFSC) for expert review of  
 this work; Valéria Ferreira Lopes, Diogo Henrique Costa de Rezende and  
 Carlos Alberto Salvador Montoya for support with molecular analysis and field  
 trips; the curators of the herbaria mentioned in this work for giving the authors  
 access to the collections; those that manage the areas where collections were  
 made. This research was partially supported by  
 FUNPESQUISA/PROPESQ/UFSC (edital N° 004/2010); CNPq (“Identificação  
 Molecular de Fungos do Brasil” MCT/CNPq/FNDCT N°. 50/2010 -  
 564944/2010-6); CAPES (“A biologia molecular como ferramenta para  
 taxonomia de fungos e de organismos fotossintetizantes”, PNPD2011 –

23038.007790/ 2011-93). CAPES provided Masters Scholarship to MABS. This work is part of project “Fungos poliporoides (Agaricomycetes) em Santa Catarina” (BOT/UFSC nº 2011.0182).

2. Laboratorio de Micología, Instituto Multidisciplinario de Biología Vegetal, CONICET - Universidad Nacional de Córdoba, CC 495, CP 5000 Córdoba, Argentina.

3. Laboratório de Pesquisa em Microbiologia (LAPEM), Departamento de Ciências Biológicas, Universidade Estadual de Feira de Santana, Avenida Transnordestina, Feira de Santana, 44036-900, BA, Brazil.

## INTRODUCTION

The occurrence of cryptic species among the morphospecies concept has been documented in South America (Decock *et al.* 2007). The resolution of taxonomic complexes is as difficult as the perception that a speciation is in course. According with Decock *et al.* (2007) and Rajchenberg & Pildain (2012), cryptic species, species complexes and the process of speciation in courses should be approached with different tools (morphological, ecological and molecular).

*Phellinus iptadeniae* Teixeira (*Hymenochaetaceae* Donk) were described in 1950, as a specialist on *Piptadenia gonoacantha* (Mart.) J.F.Macbr (*Fabaceae*, as *P. communis* Benth.) from Atlantic forest of São Paulo State (Teixeira 1950). Drechsler-Santos *et al.* (2010) expanded its distribution to the Caatinga biome, reporting it as host-recurrent on *Piptadenia spp.* In the Atlantic Rain Forest of Southeast Brazil, state of Santa Catarina, *Phellinus grenadensis* (Murrill) Ryvardeen has been reported exclusively on *Piptadenia gonoacantha* (Gerber & Loguercio-Leite 2000, Drechsler-Santos & Loguercio-Leite 2006). Remarkably, these records are based on specimens that strikingly are morphologically similar to *Phellinus iptadeniae*. *Phellinus grenadensis* was originally described from the Caribbean island of Grenada, and is reported as widely distributed in the tropics growing on different plant families (Ryvardeen 2004).

The aims of this work were: (i) to confirm the presence of *P. iptadeniae* on *Piptadenia spp.*, extending the distribution to the Southeast Atlantic Rain Forest of Brazil (Santa Catarina states), and (ii) to delimit the taxon, using molecular, morphological and ecological data.

## MATERIAL AND METHODS

Fresh specimens of the presumably *Phellinus grenadensis* growing on *Piptadenia gonoacantha* in Santa Catarina state were collected, analyzed in detail (molecular, macro- and microscopically) and properly kept at Herbarium FLOR. Type material of *Phellinus iptadeniae* (IAC 4365, Figure 1cd) and *Phellinus grenadensis* (NYGB 742990, Figure 1ab) and representative materials from Caatinga biome were also studied for further morphological comparisons (Table 1, n=40). Microscopic observations were made from slide preparations stained with 3-5% KOH and Phloxine, or Melzer's reagent. Sections of the basidiomes were incubated for two days at 40°C in NaOH 3%

solution, then carefully dissected under a stereomicroscope and examined in NaOH 3% solution at room temperature (Decock *et al.* 2013). Herbaria acronyms are based on Index Herbariorum (Thiers, continuously updated).

Material examined (*Phellinus piptadeniae*): **Brazil**. São Paulo: Campinas, Bosque dos Jequitibás, 12.IX.1943, *A.R.Teixeira e P.R.Santos* (IAC 4365, holotype, figure 1cd); Santa Catarina: Florianópolis, Campus Universitário/UFSC, 25.I.2011, *Borba-Silva MABS 106* (FLOR 39571); *ibid*, *Borba-Silva MABS 107* (FLOR 39572), 14.IV.2011, *MABS 135* (FLOR 39573); *ibid*, *MABS 136* (FLOR 39574, figure 1ef); *ibid*: Santo Amaro da Imperatriz, Caldas da Imperatriz, BR 282 KM 26, 31.X.2012, *Drechsler-Santos et al. DS 846* (FLOR, figure 1g); *ibid*: Palhoça, Parque Estadual da Serra do Tabuleiro, 27.IX.2000, Groposo 037 (FLOR 11895, as *Phellinus grenadensis*); *ibid*, 18.VII.2001, Groposo & Andrade 176 (FLOR 11957, as *Phellinus grenadensis*); *ibid*, 15.V.2005, *Michels et al. 509* (FLOR 31816, as *Phellinus grenadensis*); Pernambuco: Caruaru, Estação Experimental do IPA (8°13'53" S x 35°55'12" W, 562m alt.), 10.XII.2008, *Drechsler-Santos et al. DS109PE* (URM80322, figure 1h); *ibid*: Triunfo (7°52'53"S x 38°06'15" W, 550-800m alt.), 26.11.2009, *Robledo 1982* (CORD), on *Piptadenia* sp.

Additional material examined (*Phellinus grenadensis*): **Grenada**. Saint George: Annandale Mountain Forests, II.1906, *W.E.Broadway* s/n (NYGB 742990, holotype, figure 1ab).

Molecular studies (Table1) have been carried out in the laboratories of Universidade Federal de Santa Catarina (gDNA extraction and target region amplification) and Universidade Estadual de Feira de Santana (sequencing). The method developed by Doyle and Doyle (1987), modified by Góes-Neto *et al.* (2005), were used for DNA extraction. The nrITS region (fungal DNA barcode region) of the extracted DNA was amplified using the universal primers ITS4 and ITS5 (Schoch *et al.* 2012). Besides the new sequences generated, we were also used ITS sequences from Caatinga specimens of *Phellinus piptadeniae* (unpublished data by Drechsler-Santos *et al.*) and related species/genera of *Hymenochaetaceae* from Genbank to construct the phylogenetic trees. GenBank accession numbers of the sequences obtained are indicated in the tree (Figure 3) after the taxa names. DNA sequences were aligned on MEGA version 5.0 (Tamura *et al.* 2011) and then edited with BioEdit version 7.1.3.0 (Hall 1999). Phylogenetic trees generated by MEGA were based on maximum parsimony (MP) and Neighbor-Joining (NJ). Bootstrap analysis (1000 bootstrap replicates)

was used to determine branch support. *Trichaptum abietinum* (Dicks.) Ryvar den was used as out group (GenBank Fj608591.1). The GeneBank access numbers of new sequences are not mentioned because the data was not published yet.

Table 1. Information of partial ITS rDNA sequences used in the phylogenetic analysis

Species	Accession number	Voucher specimens	Origin	Hosts
<i>Phellinusiptadeniae</i>	unpublished	FLOR39571	Brazil	<i>Piptadenia gonoacantha</i>
<i>Phellinusiptadeniae</i>	unpublished	FLOR39573	Brazil	<i>Piptadenia gonoacantha</i>
<i>Phellinusiptadeniae</i>	unpublished	FLOR39574	Brazil	<i>Piptadenia gonoacantha</i>
<i>Phellinusiptadeniae</i>	unpublished	URM80322	Brazil	<i>Mimosa sp.</i>
<i>Phellinusiptadeniae</i>	unpublished	URM80345	Brazil	<i>Senegalia sp.</i>
<i>Phellinusiptadeniae</i>	unpublished	URM80361	Brazil	<i>Senegalia sp.</i>
<i>Phellinus sp.</i>	unpublished	URM80362	Brazil	not informed
<i>Phellinusfastuosus</i>	AY558615	not informed	not informed	not informed
<i>Phellinusrobiniae</i>	AY558646.1	CBS211.36	not informed	not informed
<i>Phellinusignarius</i>	GQ383717	not informed	not informed	not informed
<i>Phellinusalni</i>	GQ383775	BRNM714864	Finland	<i>Alnus incana</i>
<i>Inonotushispidus</i>	JX501315	B61	not informed	not informed
<b>Outgroup</b>				
<i>Trichaptum abietinum</i>	FJ7686761	1302BG	not informed	not informed

## RESULTS AND DISCUSSION

The morphological analysis (Table 2) showed that the characters: hymenophore (thickness dissepiment, diameter and number of pores per mm) and basidiospores (size and shape) were not diagnostic at species level in this group, *i.e.*, there are no differences among the specimens studied. However, the type of hyphal system, black line in the context and the length of skeletal hyphae of the tubes were characters that distinguish *Phellinus grenadensis* from the other specimens studied, previously named as *P. piptadeniae*. *Phellinus grenadensis* is dimitic in the tubes and in the context, while *P. piptadeniae* is dimitic in the tubes and monomitic in the context. Additionally, *P. piptadeniae* presents very distinct black line (figure 1dfh) in the context and longer skeletal hyphae (figure 2ab) than *P. grenadensis*. Furthermore, the specimens from Atlantic Rain Forest of Santa Catarina (ARF of SC) and the holotype of *P. piptadeniae* were collected exclusively in *Piptadenia gonoacantha*, while collections of *P. grenadensis* were recorded widely distributed to the America and Africa on many other hosts (Carranza and Ryvardeen 1998, Ryvardeen and Guzmán 1993, Dennis 1970, Gilbertson and Ryvardeen 1987, Gilbertson *et al.* 2002). In this case (table 1), the data clearly show that the specimens collected in the ARF of SC are not *P. grenadensis*, and consequently, correspond to the species *P. piptadeniae*.

**Table 2.** Morphological, host and geographical distribution data of *Phellinus grenadensis* and *Phellinus piptadeniae*.

Specimens	<i>P. grenadensis</i> Type	<i>P. piptadeniae</i> Type	<i>P. piptadeniae</i> FLOR 39571*	<i>P. piptadeniae</i> FLOR 39573*	<i>P. piptadeniae</i> FLOR 39574*	<i>P. piptadeniae</i> (CORD)	<i>P. piptadeniae</i> (URM80322)
Local specimen (geographical distribution)	Grenada Island (Africa and Americas)	Campinas (Brazil: ARF of SP)	SC Island (Brazil: ARF of SC)	SC Island (Brazil: ARF of SC)	SC Island (Brazil: ARF of SC)	Triunfo (Brazil: Caa of PE)	Caruaru (Brazil: Caa of PE)
Host species	Unknown	<i>P.gonoacantha</i>	<i>P.gonoacantha</i>	<i>P.gonoacantha</i>	<i>P.gonoacantha</i>	<i>Piptadenia sp.</i>	<i>Mimosa sp.</i>
Pores/mm (average)	5-7 (6.2)	3-5 (4.2)	5-8 (6.0)	5-8 (5.8)	5-8 (5.8)	4-6 (5.1)	4-6 (4.7)
Pores diameter (average) in $\mu\text{m}$	125-166 (145.4)	130-200 (176.8)	130-280 (204.5)	140-200 (167.1)	145-200 (171)	100-200 (150)	160-230 (188.8)
Dissepiment thickness (average) in $\mu\text{m}$	40-110 (68.4)	30-150 (126.5)	30-110 (66.9)	35-111 (67.3)	29,5-130,5 (72.8)	50-140 (84.3)	60-160,5 (101.3)
Context (black line present)	indistinct	distinct	distinct	distinct	distinct	distinct	distinct
Hyphal system structure (mitism)	Di	Mono/Di	Mono/Di	Mono/Di	Mono/Di	Mono/Di	Mono/Di
Skeletal hyphae of tubes Length range (mean) in $\mu\text{m}$	99,3 - 331 (204)	300 - 482 (391)	280 - 450,5 (354)	224,5 - 461,3 (346)	224,2 - 552,1 (356)	309 - 486,5 (373)	338 - 630 (409)
Size of spores (mean) in $\mu\text{m}$	4,5-5,5 x 3,5- 4,5 (4,9 x 4,1)	4,5-6,0 x 3,5-5,0 (5,3 x 4,2)	4,0-5,0 x 3,0-4,5 (4,6 x 3,8)	5,0-5,5 x 4,0-4,5 (5,2 x 4,2)	5,0-6,0 x 4,0-4,5 (5,6 x 4,3)	4,5-6,0 x 3,5-4,5 (5,4 x 4,0)	5-5,5 x 4-4,5 (5,1 x 4,1)
Spore Quotient (Q) in $\mu\text{m}$ and shape	1,20 - BE	1,28 - BE	1,20 - BE	1,24 - BE	1,29 - BE	1,29 - BE	1,25 - BE

\* Specimens included in the phylogenetic analysis. ARF = Atlantic Rain Forest; BE = broadly ellipsoid ( $Q = 1,15 - 1,30 \mu\text{m}$ , Largent et al. 1977); Caa = Caatinga; Di = Dimitic; Mono/Di = Monomitic in the context and Dimitic in the tubes; PE = Pernambuco states; SP = São Paulo states; SC = Santa Catarina states.



**Figure 1.** Basidiomes of *Phellinus grenadensis* (a-b) and *P. piptadeniae* (c-h): a. upper surface of holotype; b. tubes layers with context among them of holotype; c. upper surface of holotype; d. context with black line and tubes layers of holotype; e-f. upper surface and context with black line and tubes layers of MABS 136 (FLOR 39574); g. specimens on *Piptadenia gonoacantha* from ARF of SC; h. context with black line and tubes layer of DS109PE (URM80322). (Scale bar = a-d: 5 cm; e-f: 10 cm; g: 20cm; h: 1cm).



According to the macro and micro morphological characters presented in the Table 1, there are no significant differences among the specimens from Atlantic Rain Forest of SC state, previously named as *P. grenadensis*, the type of *P. piptadeniae* from Atlantic Rain Forest of SP state, and those specimens from Caatinga biome in PE state. The exception was the fact that they are from different regions and were founded on different host species (Table 1). The number of pores/mm of holotype of *P. piptadeniae* (3-5) seems to be distinct with is compared with those of the Atlantic Rain Forest of SC and Caatinga of PE specimens (5-8 and 4-6, respectively). However, when considering the pore diameter and the dissepiment thickness there are not differences. On the other hand, the specimens of *P. piptadeniae* from Atlantic Rain Forest of SC, including the type material from SP, present bigger basidiomes with the pilear surface indurating that becomes rimose and more stratified tube layers than those from Caatinga of PE (figure 1fh). These differences could be understood as only morphological variations of different populations of the same species, or as different stages of ontogenetic development of that species. Then, morphological and ecological (host and geographic distribution) data suggested *P. piptadeniae* as a taxonomic complex.

The occurrence of cryptic species among the morphospecies concepts seems to be commonly founded when one studies fungal taxa nowadays (Decock *et al.* 2007). For instance, Fischer & Binder (2004), when trying to resolve the relationships among 69 collections of the genera *Phellinus* s.s. Quél. and *Fomitiporia* Murrill (*Hymenochaetaeaceae*, *Hymenochaetales*), from the Northern Hemisphere, pointed out that the sole application of morphological criteria was of limited value especially in taxa exhibiting a wide distribution range, both in terms of geographic origin and ecological niche. The authors, using an integrative approach (morphological, phylogenetic and biological species recognition) detected 12 species among the seven species previously recognized by the application of a strict morphological approach.

Another possible challenge when working with taxonomic complexes is faced with a speciation processes. Rajchenberg & Pildain (2012), from a detailed studies and comparisons of brown-rot polypore *Ryvardenia cretacea* (Lloyd) Rajchenb. (*Polyporaceae*, *Polyporales*), did not show the existence of significant morphological differences between populations from Australia (Tasmania, Victoria) and Argentina (Patagonia), although the phylogenetic analysis (nrITS) revealed the existence of two well supported clades. The results of this study support

the idea that genotypic differentiation precedes phenotypic and reproductive isolation and provide further evidence on the complex nature of issues regarding species concepts, recognition, and delimitation (Urcelay *et al.* 2012).

Both aforementioned studies, Fischer & Binder (2004) and Rajchenberg & Pildain (2012), used the nuclear ribosomal ITS region as a complementary tool to recognize cryptic species or cryptical speciation, respectively. Furthermore, the ITS region is considered nowadays as a universal DNA barcode marker for fungi (Schoch *et al.* 2012).

Our phylogenetic analysis of the nrITS region include 873 total characters, of which 514 were parsimony informative. Bootstrap values obtained using MP and NJ were very similar and both analyses resulted in a similar tree topology (the consensus tree is presented in figure 3), with high statistical support for the *Phellinus piptadeniae* complex (MP = 100 and NJ = 99). Geographical/ecological sub-groups within *P. piptadeniae* are well supported with high bootstrap values for Caatinga specimens of PE (MP = 93 and NJ = 97) and for ARF specimens of SC (MP = 99 and NJ = 99).


Although, the Caatinga and AFR Clades are inclusive of a single branch (*Phellinus piptadeniae*), they are both highly supported, evidenced that the Caatinga and ARF specimens should be a different species, according to the phylogenetic species concept. These two populations may be the result of their own evolutionary history with their hosts, inferring in this case, that the isolation has occurred according to the specialization with the hosts in particular. As observed, the Atlantic Rain Forest populations are

**Figure 2.** Skeletal hyphae: a. *Phellinus grenadensis* (holotype); b. *Phellinus piptadeniae* (FLOR 39573) (scale barr = 50  $\mu$ m).



specific on *Piptadenia gonoacantha* and, although Drechsler-Santos *et al.* (2010) mentioned the recurrence of *P. piptadeniae* on *Piptadenia* spp. as hosts in the Caatinga of National Park of Catimbau (Pernambuco states), the Caatinga population studied here were founded on other genera hosts (*Senegalia* sp. and *Mimosa* sp., Figure 3).

**Figure 3.** Phylogenetic relationships of *Phellinus piptadeniae* complex (indicated by bars, including the inclusive Caatinga and ARF clades and their hosts) and other species and genera [*Inonotus hispidus* (Bull.) P. Karst., *Phellinus fastuosus* (Lév.) Ryvarden, *P. robiniae* (Murrill) A. Ames, *P. igniarius* (L.) Qué!., *P. alni* (Bondartsev) Parmasto] of *Hymenochaetaceae*, based on complete sequence of nrDNA ITS. The tree above was generated by heuristic search using Maximum Parsimony (MP) and Neighbor-Joinin (NJ) and the Bootstrap values (1000 replicates) are indicated at the internal nodes, respectively. The sequences of the materials collected in ARF of SC are represented by the numbers of collector (MABS) and those from Caatinga biome of PE are represented by the numbers of collector (Drechsler-Santos, unpublished data by Drechsler-Santos *et al.*). The sequences obtained from GenBank are labeled with the corresponding accession number, using *Trichaptum abietinum* as an outgroup.



As suggested by Hallenberg et al. (2007), geographically delimited populations of a fungal species may often be distinguished phylogenetically. Accordingly, *Phellinus piptadeniae* presents ecologically (*i.e.* geographical and/or host distribution) and phylogenetically two well delimited populations represented by the two inclusive clades (Caatinga and ARF of SC) in the Figure 3. However, according to Rajchenberg & Pildain (2012), when no morphological differences exist and no compatibility tests were made between segregated populations it is better to understand them as species complexes. For the time being our conclusion is that *Phellinus piptadeniae* represent a taxonomic complex of species that is undergoing a speciation process as evidenced by the existence of two phylogenetic lineages, *i.e.*, two ecological (host and geographic distribution) populations. This work is the first step to assess the relationships within *Phellinus piptadeniae*. Further research including compatibility tests (biological species concept) and more sample sequences of different hosts, as well as multigenic phylogenetic studies, are necessary to further understand the taxonomy and ecology within this complex.

## LITERATURE CITED

- Carranza, J. and L. Ryvardeen. 1998. Additional list of pore fungi of Costa Rica. *Mycotaxon* 69: 377-390.
- Decock, C. M. Amalfi, G. Robledo, G. Castillo. 2013. *Phylloporia nouraguensis*, an undescribed species on Myrtaceae from French Guiana. *Cryptogamie Mycologie* 34 (1): 15-27.
- Decock, C., S.H. Figueroa, G. Robledo, G. Castillo. 2007. *Fomitiporia punctata* (Basidiomycota, Hymenochaetales) and its presumed taxonomic synonyms in America: taxonomy and phylogeny of some species from tropical/subtropical areas. *Mycologia* 99(5): 733-52.
- Dennis, R.W.G. 1970. Fungus flora of Venezuela and adjacent countries. *Kew Bulletin Additional Series* 3: i-xxxiv, 531 p.
- Doyle, J.J. and J.L. Doyle. 1987. A rapid isolation procedure for small quantities of fresh tissue. *Phytochemical Bulletin* 19: 11-15.
- Drechsler-Santos, E.R. and C. Loguercio-Leite. 2006. Basidiomycetos SC – Brasil. Guia Eletrônico de Identificação. Version BASC – 1.0. Acesso: <http://www.ccb.ufsc.br/bot/micologia/> em 20/06/2011.
- Drechsler-Santos, E.R., P.J.P.Santos, T.B. Gibertoni, M.A.Q.Cavalcanti. 2010. Ecological aspects of Hymenochaetales in an area of Caatinga (semi-arid) in Northeast Brazil. *Fungal Diversity* 42: 71-78.
- Fisher, M. and M. Binder. 2004. Species recognition, geographic distribution and host-pathogen relationships: a case study in a group of lignicolous basidiomycetes, *Phellinus* s.l. *Mycologia* 96 (4): 799-811.
- Gerber, A.L. and C. Loguercio-Leite. 2000. Polyporoid Wood-Rotting Fungi (Basidiomycetes) II – New Records from Southern Brazil. *Mycotaxon* 76: 175-185.
- Gilbertson, R.L. and L. Ryvardeen. 1987. North American Polypores Vol. 2. *Megasporoporia - Wrightoporia*. *Fungiflora*, Oslo, 434-885 p.

Gilbertson, R.L., D.M. Bigelow, D.E. HEMmes, D.E. Desjardin. 2002. Annotated check list of Wood-Rotting Basidiomycetes of Hawaii. *Mycotaxon* 82: 215-239.

Góes-Neto, A. and C. Loguercio-Leite , R.T. Guerrero. 2005. DNA extraction from frozen field-collected and dehydrated herbarium fungal basidiomata: performance of SDS and CTAB-based methods. *Biotemas* 18(2): 19-32.

Hall, T.A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95-98.

Hallenberg N., R.H. Nilsson, A. Antonelli, S.-H. Wu, N. Maekawa, B. Norden. 2007. The *Peniophorella praetermissa* species complex (Basidiomycota). *Mycological Research* 111: 1366-1376.

Largent et , D., D. Johnson, R. Watling. 1977. How to Identify Mushrooms to Genus III: Microscopy Features. Mad River Press Inc. California. 122 p.

Rajchenberg, M. and M.B. Pildain. 2012. Molecular studies reveal a speciation process within *Ryvardenia cretacea* (Polyporales, Basidiomycota). *Kurtziana* 37(1): 7-13.

Ryvarden, L. and G. Guzmán. 1993. New and interesting polypores from Mexico. *Mycotaxon* 47: 1-23.

Schoch, C. L., K. A. Seifert., S. Huhndorf, V. Robert, J. L. Spouge, C. A. Levesque, W. Chen and Fungal Barcoding Consortium. 2012. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *PNAS* (109)16: 6241-6246.

Tamura, K., D. Peterson, N. Peterson, G. Stecher, M. Nei and S. Kumar. (2011) MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Molecular Biology and Evolution* 28: 2731-2739. doi:10.1093/molbev/msr121.

Teixeira, A.R. 1950. Himenomicetos Brasileiros V. *Bragantia* 10(4): 113-134.



Thiers, B. [continuously updated]. Index Herbariorum: a global directory of public herbaria and associated staff. New York Garden's Virtual Herbarium. <<http://sweetgum.nybg.org/ih/>>.

Urcelay, C. and G. Robledo. 2004. Community structure of polypores (Basidiomycota) in Andean alder wood in Argentina : Functional groups among wood-decay fungi? *Austral Ecology* 29: 471-476.

Urcelay, C. and G. Robledo. 2009. Positive relationship between wood size and basidiocarp production of polypore fungi in *Alnus acuminata* forest. *Fungal Ecology* 2(3): 135-139.

Urcelay, C., A. Romero and G. Robledo. 2012. Taxonomy and Systematics of fungi from South America – a special issue in honoring Leif Ryvarden: Introduction . *Kurtziana* 37(1): 3-6.

## CONSIDERAÇÕES FINAIS

O estudo realizado mostrou que os fungos políporoides da Mata Atlântica de Santa Catarina, podem ser agrupados funcionalmente pela sua frequência relativa no tipo de substrato em que ocorrem. Comparado ao estudo de grupos funcionais realizado na região das Yungas argentinas, os grupos funcionais são semelhantes, porém compostos por espécies distintas. *Fuscoporia gilva*, *Pycnoporus sanguineus*, *Polyporus tricholoma* e *Trametes villosa* ocorrem nos diferentes ecossistemas, Mata Atlântica e Yungas, e nos mesmos grupos funcionais, com exceção da primeira, que além de ser registrada em galhos mortos também foi encontrada em galhos vivos nas Yungas.

Com relação à riqueza e abundância de espécies, *Polyporus dictyopus*, *Perenniporia martia* e *Fuscoporia wahlbergii*, mostraram-se como espécies dominantes para cada um dos três grupos funcionais saprófitas, diferentemente das espécies dominantes registradas na Yungas argentinas.

O conhecimento dos políporos da área estudada e da sua relação direta com os tipos de substratos, podem facilitar a elaboração de estratégias de conservação deste bioma em particular, uma vez que a abundância e a riqueza de espécies encontradas podem estar relacionadas com a oferta dos substratos.

Os registros de espécimes himenoquetoides coletadas sobre *Piptadenia gonoacantha* em Santa Catarina, tradicionalmente identificados como *Phellinus grenadensis*, foram confirmados como *Phellinus piptadeniae* a partir das análises morfológicas.

A utilização de uma abordagem integrativa, *i.e.*, a partir de análises morfológicas, filogenéticas e ecológicas (hospedeiro e distribuição geográfica), se mostrou eficiente no entendimento do complexo taxonômico estudado. Obviamente, testes de compatibilidade (conceito biológico) são sugeridos para a resolução de *Phellinus piptadeniae complex*.